

Post-treatment	S1		S2		S3		S4		S5		S6		S7		S8	
24hr	L	R	L	R	L	R	L	R	L	R	L	R	L	R	L	R
1	1.6	1.4	1.2	1.5	1.3	1.7	1.3	1.6	0.9	1.6	1.4	1	1.6	1.2	1.6	1.9
2	1.6	1.1	1.1	1.4	1.2	1.5	1.2	0.9	1.1	0.8	1.2	1.5	1	1.3	1.2	1
3	1.5	1	1	1.3	1.1	1.4	1.1	0.9	1.4	1.6	0.9	0.8	0.7	0.5	1.9	0.9
4	1.3	0.7	1	1.2	0.7	1.1	0.9	0.5	1.7	0.8	0.9	1	1.3	1.2	1	1.4
5	0.8	0.6	0.9	1.1	0.5	1.1	0.8	0.5	0.7	0.8	0.7	0.5	1.2	1.6	0.8	0.9
average	1.4	1.0	1.0	1.3	1.0	1.4	1.1	0.9	1.2	1.1	1.0	1.0	1.2	1.2	1.3	1.2
	ipsilateral		controlateral													
average	1.13	1.12														
SD	0.14	0.17														
SE	0.05	0.06														

Post-treatment	CFA1		CFA2		CFA3		CFA4		CFA5		CFA6		CFA7		CFA8	
24hr	L	R	L	R	L	R	L	R	L	R	L	R	L	R	L	R
1	0.7	1.3	0.1	2.4	0.5	1.4	0.9	1.3	0.1	0.5	0.1	1	0.1	0.6	0.2	1.4
2	0.5	1.2	0.1	2.4	0.5	1.3	0.9	1.1	0.3	1	0.1	1.4	0.3	0.6	0.4	0.9
3	0.5	1.2	0.1	2.2	0.5	1	0.7	0.9	0.1	0.6	0.2	1	0.1	0.8	0.1	0.8
4	0.5	0.9	0.1	1.7	0.1	0.9	0.6	0.7	0.2	1.1	0.2	0.8	0.3	1.2	0.5	1.2
5	0.3	0.9	0.1	1.2	0.1	0.8	0.4	0.7	0.4	0.8	0.4	0.6	0.2	1.3	0.5	1.1
average	0.5	1.1	0.1	2.0	0.3	1.1	0.7	0.9	0.2	0.8	0.2	1.0	0.2	0.9	0.3	1.1
	ipsilateral		controlateral													
average	0.33	1.11														
SD	0.19	0.37														
SE	0.07	0.13														

0.9% NaCl	ipsilateral	controlateral	CFA	ipsilateral	controlateral
average	1.13	1.12	average	0.33	1.11
SD	0.14	0.17	SD	0.19	0.37
SE	0.05	0.06	SE	0.07	0.13

ANOVA	p value
CFA-cont vs. CFA-ipsi	<0.01
0.9% NaCl-ipsi vs. CFA-i;	<0.01

